

## Corporate Medical Policy

### Human Immunodeficiency Virus – M2116

**File Name:** Human\_immunodeficiency\_virus  
**Origination:** 1/2019  
**Last Review:** 5/2023

#### Description of Procedure or Service

---

##### Description

Human immunodeficiency virus (HIV) is an RNA retrovirus that infects human immune cells, specifically CD4 cells, causing progressive deterioration of the immune system ultimately leading to acquired immune deficiency syndrome (AIDS) characterized by susceptibility to opportunistic infections and HIV-related cancers (CDC, 2014). HIV-1 is the dominant subtype of HIV infection, but another subtype, HIV-2, is a crucial subtype in certain areas of the world, such as Western Africa (Wood, 2021).

##### Scientific Background

Human immunodeficiency virus (HIV) targets the immune system, eventually hindering the body's ability to fight infections and diseases. If not treated, an HIV infection may lead to acquired immunodeficiency syndrome (AIDS) which is a condition caused by the virus. There are two main types of HIV: HIV-1 and HIV-2; both are genetically different. HIV-1 is more common and widespread than HIV-2.

##### *HIV-1*

Human immunodeficiency virus type 1 (HIV-1) RNA in blood can be measured using qualitative or quantitative techniques. Qualitative testing is used as a screening test to identify HIV-infected individuals whereas quantitative measurement of HIV-1 viral loads in the blood is used in management and monitoring of HIV-1 infected individuals. HIV-1 RNA levels may also be used to establish the diagnosis of HIV infection in specific situations where combination tests that detect HIV p24 antigen and HIV antibodies are not appropriate (neonatal or acute infection) (Caliendo, 2022).

Three primary real-time reverse transcriptase polymerase chain reaction (RT-PCR) commercial tests are commonly used to quantify HIV-1 RNA from plasma. These tests are more sensitive (detecting 20 to 40 copies/mL of HIV RNA), have a broader linear range (detecting virus to at least 10 million copies/mL), and pose a lower risk of carry over contamination than prior PCR assays. The tests are “COBAS TaqMan HIV-1 Test version 2” by Roche Diagnostics, “RealTime HIV-1” by Abbott Molecular, and “Aptima HIV-1 Quant Dx Assay” by Hologic (Caliendo, 2022). In 2020, the Aptima assay received FDA approval to aid in diagnosis, in addition to its original use of quantitation (BusinessWire, 2020; FDA, 2020).

Sources of variability between HIV-1 assays include differences in technology platform, plasma input volume, and ability to detect HIV-1 subtypes. Monitoring of individual patients should be performed on the same technology platform to ensure appropriate interpretation of changes in viral load (Sollis et al., 2014). An important difference between assays is the gene target; with the

# Human Immunodeficiency Virus – M2116

increasing use of integrase inhibitors, monitoring for resistance mutations in the integrase gene is essential to ensure that the primer and probe binding sites are not impacted (Caliendo, 2022).

Overall, studies of real-time RT-PCR tests have shown high concordance, high correlation values, and good agreement among all assays (Mor et al., 2015). However, their manufacturers have reported that variation and error tend to increase at the lower limits of quantitation of the assays (Swenson et al., 2014). The high variability around the threshold of detectability of the viral load assays should be noted since many patients have viral loads in this range. Agreement between these assays was improved using a 200-copies/ml threshold (Swenson et al., 2014) consistent with the current HIV treatment guidelines' definition of virological failure (Saag et al., 2020).

Furthermore, changes in HIV-1 RNA levels must exceed at least 0.5 log<sub>10</sub> or threefold in magnitude to represent biologically relevant changes in viral replication (Hughes et al., 1997; Saag et al., 1996). Viral RNA levels can also transiently rise due to acute illness, herpes outbreak, or vaccination; however, values usually return to baseline within one month (Caliendo, 2022). CD4 cell counts are weakly correlated with viral RNA measurements. Viral RNA measurements, although, do not replace CD4 cell counts in the management of HIV-1-infected patients and should be used in parallel (Caliendo, 2022).

## *HIV-2*

Human immunodeficiency virus type 1 (HIV-2) is another subtype of HIV. Compared to HIV-1, HIV-2 appears milder clinically; it is characterized by a longer asymptomatic stage, slower declines of CD4 cell counts, and lower levels of plasma viremia in chronically ill patients (Gottlieb, 2022b). However, these numerical thresholds are not as well-defined as those of HIV-1 as there is currently not as much data available for HIV-2. Further, although quantification of HIV-2 RNA viral load may be useful, it is not widely commercially available, as the few labs that offer HIV-2 testing only offer qualitative testing and not quantitative (Gottlieb, 2022a). This is particularly crucial as HIV-1 assays typically do not properly detect HIV-2 viral load (DHHS, 2022a). It is possible for commercially available HIV-1 diagnostic assays to cross-react with HIV-2, disrupting the results. A reactive HIV-1 Western Blot may not be indicative of a true HIV-1 infection. For example, a patient may have reactive HIV serology, but test negative on a confirmatory HIV-1 Western blot. This scenario may indicate an HIV-2 infection. Clinical manifestations of HIV-2 infection are generally similar to HIV-1 infection, but much remains to be discovered about the general course of HIV-2 infection (Gottlieb, 2022a).

Despite HIV-2's milder symptoms, certain clinical features may make an infection more difficult to manage; for example, HIV-2 is intrinsically resistant to non-nucleoside reverse transcriptase inhibitors, as well as enfuvirtide. Assessment of genotypic or phenotypic resistance is also unexplored, with no currently FDA-approved genotypic or phenotypic resistance assays available (DHHS, 2022a).

Although HIV-2 is endemic to West Africa (Senegal, Gambia, Guinea-Bissau, et al.) the epidemiological trends may be shifting; the CDC only reported 166 cases of HIV-2 from 1987 to 2009 but this may be underestimated as HIV-2 is often asymptomatic. There were 24 cases of HIV-2 identified in New York City between 2010 and 2020, with 25 additional probable cases. Additionally, as much as 5% of HIV cases are thought to be HIV-2 (Gottlieb, 2022b; Quinn, 2022).

## *Drug Resistance*

HIV replicates rapidly; a replication cycle rate of approximately one to two days ensures that after a single year, the virus in an infected individual may be 200 to 300 generations removed from the initial infection-causing virus (Coffin & Swanstrom, 2013). This leads to great genetic diversity of each HIV infection in a single individual. As an RNA retrovirus, HIV requires the use of a reverse transcriptase for replication purposes. A reverse transcriptase is an enzyme which generates

# Human Immunodeficiency Virus – M2116

complimentary DNA from an RNA template. This enzyme is error-prone with the overall single-step point mutation rate reaching about  $3.4 \times 10^{-5}$  mutations per base per replication cycle (Mansky & Temin, 1995), leading to approximately one genome in three containing a mutation after each round of replication (some of which confer drug resistance). This rate is comparable to other RNA viruses. This pace of replication, duration of infection, and size of the replicating population allows the retrovirus to evolve rapidly in response to selective influences (Coffin & Swanstrom, 2013).

Due to the high rate of mutation in HIV viruses, drug resistance mutations are common. Some drugs may be resisted by a single mutation—these drugs have a “low genetic barrier” to resistance. Such mutations are common enough to be termed “signature mutations,” which are frequently associated with a specific drug resistance. For example, the K103N mutation commonly leads to resistance for efavirenz. Efavirenz is a standard retroviral medication used to treat and prevent HIV and AIDs. To combat this, medical professionals can now assess drug-resistant HIV variants using phenotypic testing and genotypic testing (Kozal, 2019a).

Genotypic assays detect the presence of specific drug-resistance mutations in several different genes (protease, reverse transcriptase, and integrase genes). For example, assays may test for resistance in nucleoside reverse transcriptase inhibitors (NRTIs), non-nucleoside reverse transcriptase inhibitors (NNRTIs), or protease inhibitors (PIs). The definition of a resistance conferring mutation is blurred, but generally includes one or more of the following conditions:

- The mutation confers phenotypic resistance when introduced into a drug-sensitive laboratory strain of HIV.
- The mutation is selected for during serial in vitro passage of the virus in the presence of a drug.
- The mutation is selected for during clinical therapy with that drug.
- The presence of the mutation in clinical isolates is associated with phenotypic resistance and virologic failure (Kozal, 2019b).

Interpretation of genotypic data may be done either by clinical expertise or through a database (in which the genotype is correlated with the phenotype). Phenotypic resistance assays measure the extent to which an antiretroviral drug inhibits viral replication. Phenotypic testing typically assesses the fold-change in susceptibility of a patient’s virus and the treatment response, while also correlating the mutations present with the fold-change in susceptibility. Recombinant virus assays (RVAs) are used; protease, reverse transcriptase, or integrase gene sequences from circulating viruses are inserted into a reference strain of HIV, and this new HIV strain is measured by the phenotypic assay (Kozal, 2019b).

Several HIV genotypic assays are available. The ViroSeq HIV-1 Genotyping system by Abbott helps to detect HIV-1 genomic mutations that may lead to resistance to certain types of antiretroviral drugs (ThermoFisher, 2011). The ATCC® HIV-1 Drug Resistance Genotyping Kit has been developed by the American Type Culture Collection (ATCC), the Centers for Disease Control and Prevention (CDC) and Thermo Fischer Scientific; this is a real time- polymerase chain reaction (rt-PCR) assay which may help to identify and monitor HIV-1 drug resistance (ATCC, 2014).

The primary phenotypic assay is “PhenoSense” from LabCorp. The Human Immunodeficiency Virus 1 (HIV-1) PhenoSense GT® Plus Integrase (Monogram® Phenotype + Genotype) test by LabCorp measures HIV genotypic and phenotypic resistance from plasma samples (LabCorp, 2021).

Advantages of the genotype assays include lower cost and shorter turnaround time. However, interpretation of these assays is complicated by combinations of individual mutations that may have a differential effect on resistance that differs from the individual mutation alone (Kozal, 2019b). Mutation combinations are known to cause resistance to certain drugs, but increase

# Human Immunodeficiency Virus – M2116

susceptibility to others, impact viral fitness, and contribute to major pathways of resistance; additionally, the interactions of mutations affecting various mechanisms can be difficult to predict. Over 20 rules-based genotypic interpretation systems (GIS) have been proposed (Fox et al., 2007; Kozal, 2019b).

Advantages of phenotypic assays include an ability to measure resistance more directly and examine the relative effect of multiple mutations on drug resistance. Limitations of the phenotypic assays include a longer turnaround time, greater expense, and biologic cut-offs above achievable drug levels. Phenotypic resistance assays may be helpful when evaluating HIV strains with known or suspected complex drug resistance mutation patterns as their actual resistance may not be accurately predicted by simply detecting the presence of multiple mutations (Kozal, 2019b). Both assays are limited by decreased sensitivity for low-level minority variants that comprise less than 5 to 20 percent of the virus population (Kozal, 2019b).

## *Analytical Validity*

Rosemary et al. (2018) performed a comparison of two genotyping assays, ViroSeq and ATCC (manufactured by Thermo-Fisher Scientific) kit. A total of 183 samples with a viral load  $\geq 1000$  copies/mL were sequenced by ViroSeq and randomly selected (85 successfully genotyped, 98 unsuccessfully genotyped). The ATCC kit also genotyped 115 of the 183 samples, and out of the 98 unsuccessfully genotyped samples, the ATCC kit was able to genotype 42. Overall, 127 of the 183 samples were genotyped. The authors noted that the sequences of the genotyped samples were 98% identical and had “similar HIVDR profiles at individual patient level” (Rosemary et al., 2018).

## *Clinical Utility and Validity*

Zhang et al. (2005) compared two phenotyping assays, Antivirogram and PhenoSense. Reverse transcriptase inhibitor susceptibility results were evaluated for 202 isolates from Antivirogram and 126 from PhenoSense. The authors found the median deviance for wild-type and mutant isolates to be lower for PhenoSense compared to Antivirogram, and PhenoSense was more likely to detect resistance to abacavir, didanosine, and stavudine when common drug resistance mutations were present (Zhang et al., 2005).

Hopkins et al. (2015) performed a study comparing the three main RT-PCR tests available, Aptima, COBAS TaqMan (CTM), and Abbott RealTime. The assays were evaluated based on plasma samples from 191 HIV positive patients as well as WHO International Standards (12-500 copies/mL). Aptima detected 141/191 (74%) of the HIV samples, CTM detected 145/191 (76%), and Abbott RealTime detected 119/191 (62%). The authors noted that precision decreased as the viral load got closer to the lower limit of quantification of 50 copies/mL (Hopkins et al., 2015).

Sempa et al. (2016) evaluated the utility of HIV-1 viral load as a prognostic indicator. A total of 489 patients were evaluated, and the viral load curves were evaluated on a linear scale and a logarithmic scale. The authors found that the viral load curve on the logarithmic scale was a statistically significant predictor of mortality, noting that each log<sub>10</sub> increase in viral load corresponded to a 1.63 times higher risk of mortality. However, the authors stress that the choice of variables and statistical model influences the predictive power of this metric (Sempa et al., 2016).

Shen et al. (2016) assessed the ability to predict phenotypic drug resistance from genotypic data. The authors used two machine learning algorithms to predict drug resistance to HIV protease inhibitors and reverse transcriptase inhibitors as well as the severity of that resistance from a query sequence. The accuracy of these classifications was found to be  $>0.973$  for eight PR inhibitors and  $0.986$  for ten RT inhibitors and the  $r^2$  was  $0.772$ – $0.953$  for the PR cohort and  $0.773$ – $0.995$  for the RT cohort. The algorithms' results were verified by “five-fold cross validation” on the genotype-phenotype datasets (Shen et al., 2016).

# Human Immunodeficiency Virus – M2116

Lindman et al. (2019) investigated the test performance of the Bio-Rad Geenius HIV-1/2 confirmatory assay against INNO-LIA HIV 1/2 Score and ImmunoComb HIV 1/2 BiSpot. The Geenius test is purported to differentiate between HIV-1 and HIV-2 infections. 131 samples from ART naïve HIV infected patients in Guinea-Bissau were evaluated. The Geenius test identified 62 samples as “HIV-1 reactive”, 37 as “HIV-2 reactive” and 32 as “HIV-1/2 dually reactive.” INNO-LIA identified 63 as HIV-1 reactive, 36 as HIV-2 reactive, and 32 as HIV-1/2 dually reactive. The agreement between Geenius compared to INNO-LIA and ImmunoComb was 92.4% and 84% respectively (Lindman et al., 2019).

Avram et al. (2019) compared the cost-effectiveness of measuring viral load to guide delivery in HIV-positive women and compared it to routine cesarian delivery. A theoretical cohort of 1275 women was used, and the authors produced a decision-analytic model to compare the two techniques. The average cost of a point-of-care HIV RNA viral load test was placed at \$15.22. The authors also assumed that each woman in the cohort would deliver two children. The authors defined the primary outcomes as “mother-to-child transmission, delivery mode, cesarean delivery-related complications, cost, and quality-adjusted life years”, and the cost-effectiveness threshold was \$100,000/quality-adjusted life year. The authors found that measuring viral load resulted in more HIV-infected neonates than routine cesarian delivery for all due to “viral exposure during more frequent vaginal births in this strategy.” The authors found an increased cost of \$3,883,371 and decreased quality-adjusted life years of 63 in the measurement strategy compared to the routine cesarian delivery strategy. At \$100,000/quality-adjusted life year, measuring viral load was found to be cost-effective only “when the vertical transmission rate in women with high viral load below 0.68%” (compared to a baseline of 16.8%) and “when the odds ratio of vertical transmission with routine cesarean delivery for all compared with vaginal delivery was above 0.885” (compared to a baseline of 0.3). The authors concluded that “for HIV-infected pregnant women without prenatal care, quantifying viral load to guide mode of delivery using a point-of-care test resulted in increased costs and decreased effectiveness when compared with routine cesarean delivery for all, even after including downstream complications of cesarean delivery” (Avram et al., 2019).

Raymond et al. (2020) evaluated the performance of the Vela Dx Sentosa next-generation sequencing (NGS) system for HIV-1 DNA genotypic resistance. 40 DNA samples were analyzed with Vela Dx Sentosa assay and the results were compared with Sanger sequencing. The Vela Dx Sentosa assay was 100% successful in amplifying and sequencing the protease and reverse transcriptase, and 86% successful in amplifying integrase sequences when the HIV DNA load was greater than 2.5 log copies/million cells. The Sentosa and Sanger sequencing were concordant for predicting protease-reverse transcriptase resistance in 20% of the 14/18 samples which were successfully sequenced. Sentosa was able to predict a higher level of resistance in three of the samples. The Vela Dx Sentosa predicted the prevalence of drug resistance to protease inhibitors (7%), nucleoside reverse transcriptase inhibitor (59%), nonnucleoside reverse transcriptase inhibitor (31%), and integrase inhibitors (20%). Overall, the authors conclude that the Vela Dx Sentosa assay can accurately predict HIV DNA drug resistance (Raymond et al., 2020).

Fogel et al. (2020) also analyzed the ability of next-generation sequencing methods to analyze HIV drug resistance. In this case, 145 plasma samples were analyzed using the ViroSeq HIV-1 Genotyping System and the veSEQ-HIV assay. Results were compared with the Abbott RealTime Viral Load assay. 142 HIV protease and reverse transcriptase sequences and 138 integrase sequences were obtained with ViroSeq. On the other hand, veSEQ-HIV detected 70.4% of the samples with protease, reverse transcriptase, and integrase sequences. Drug resistance mutations were detected in 33 ViroSeq samples and 42 veSEQ-HIV samples. Overall, veSEQ-HIV predicted more drug resistance mutations and worked better for larger viral loads. Results from veSEQ-HIV strongly correlated with the results from Abbott RealTime Viral Load assay. The authors conclude that the veSEQ-HIV assay provided results for most samples with higher viral loads, was accurate for predicting drug resistance mutations, but detected mutations at lower levels compared with the ViroSeq assay (Fogel et al., 2020).

# Human Immunodeficiency Virus – M2116

Pröll et al. (2022) investigated whether NGS from proviral DNA and RNA could be an alternative to using plasma viral RNA as the material of choice for genotypic resistance testing at the start of ART and virologic failure for patients with low viremia. When taking samples from 36 patients, with varying viral loads of 96 to 390,000 copies/mL, the researchers found 2476 variants/drug resistance mutations by SS, while 2892 variants were found by NGS. Researchers stated, “An average of 822/1008 variants were identified in plasma viral RNA by Sanger or NGS sequencing, 834/956 in cellular viral RNA, and 820/928 in cellular viral DNA.” This demonstrates that cellular RNA and cellular viral DNA could serve as viable substitutes when testing for variant detection and genotypic resistance among patients with HIV and low viremia (Pröll et al., 2022).

Ehret et al. (2022) tested the performance of the “Xpert® HIV-1 Viral Load (VL) XC” HIV-RNA quantitative assay made by Cepheid. This assay has been redesigned to use a dual-target approach. The authors tested 533 fresh and frozen samples from HIV-1 positive patients on the Abbott HIV-1 assay and the Xpert XC assay. “The Xpert XC assay yielded valid results in 98.5% (N = 528/536) of cases.” The authors conclude that “The Xpert XC assay showed excellent correlation with the Abbott assays for all tested HIV-1 subtypes” (Ehret et al., 2022).

## **Applicable Federal Regulations**

### **Food and Drug Administration (FDA)**

All three of the primary RT-PCR tests for HIV-1 have been approved by the FDA:

In May 2007, the FDA approved the Abbott RealTime HIV-1 Amplification Reagent Kit. From the FDA website: “The Abbott RealTime HIV-1 assay is an in vitro reverse transcription-polymerase chain reaction (RT-PCR) assay for the quantitation of Human Immunodeficiency Virus type 1 (HIV-1) on the automated m2000 System in human plasma from HIV-1 infected individuals over the range of 40 to 10,000,000 copies/mL” (FDA, 2007a).

On May 11, 2007, the FDA approved the COBAS® AmpliPrep/COBAS® TaqMan® HIV-1 Test. From the FDA website: “The COBAS AmpliPrep/COBAS TaqMan HIV-1 is an in vitro nucleic acid amplification test for the quantitation of Human Immunodeficiency Virus (HIV-1) nucleic acid in human plasma, using the COBAS AmpliPrep Instrument for automated sample preparation and the COBAS TaqMan Analyzer or COBAS TaqMan 48 Analyzer for automated amplification and detection. This test is intended for use in conjunction with clinical presentation and other laboratory markers of disease progress for the clinical management of HIV-1 infected patients” (FDA, 2007b).

In 2016, the FDA approved the Aptima® HIV-1 Quant Assay. From the FDA website: “The Aptima HIV-1 Quant assay is an in vitro nucleic acid amplification test (NAAT) for the quantitation of human immunodeficiency virus type 1 (HIV-1) RNA in human plasma from HIV-1 infected individuals on the fully automated Panther® system. The Aptima HIV-1 Quant assay quantitates HIV-1 RNA groups M, N, and O over the range of 30 to 10,000,000 copies/ mL.” On November 20, 2020, this assay was given an FDA approval for dual use for diagnosis and viral load monitoring for HIV-1 (BusinessWire, 2020; FDA, 2020).

The following screening antibody tests are FDA-approved to differentiate HIV-1 from HIV-2:

On August 26, 2019, the FDA approved the Geenius HIV-1/2 Supplemental Assay. From the FDA Website: “The Geenius™ HIV 1/2 Supplemental Assay is a single-use immunochromatographic assay for the confirmation and differentiation of individual antibodies to Human Immunodeficiency Virus Types 1 and 2 (HIV-1 and HIV-2) in serum or plasma samples (EDTA, lithium heparin, sodium citrate, and CPD) from blood donors. The Geenius™ HIV 1/2 Supplemental Assay is intended for use as an additional, more specific test for human serum and plasma samples with repeatedly reactive results by an FDA licensed blood donor screening test for

# Human Immunodeficiency Virus – M2116

antibodies to HIV-1/HIV-2. The results of the Geenius™ HIV 1/2 Supplemental Assay are read and interpreted only with the Geenius™ Reader with dedicated software.” 200 known HIV-2 positive samples were classified by Geenius, with 77 interpreted as only HIV-2 positive, 108 with HIV-2 with HIV-1 cross reactivity, 12 as undifferentiated, and 3 as HIV-2 indeterminate (FDA, 2019).

On July 23, 2015, the FDA approved the BioPlex 2200 HIV Ag-Ab Assay. From the FDA Website: “The BioPlex 2200 HIV Ag-Ab assay is a multiplex flow immunoassay intended for the simultaneous qualitative detection and differentiation of the individual analytes HIV-1 p24 antigen, HIV-1 (groups M and O) antibodies, and HIV-2 antibodies in human serum or plasma (fresh or frozen K2 EDTA, K3 EDTA, lithium heparin, sodium heparin; fresh citrate). This assay is intended as an aid in the diagnosis of infection with HIV-1 and/or HIV-2, including acute (primary) HIV-1 infection. The assay may also be used as an aid in the diagnosis of infection with HIV-1 and/or HIV-2 in pediatric subjects as young as two years of age, and pregnant women.” The test was found to differentiate all 1363 HIV-1 samples correctly and 188 of 200 HIV-2 samples correctly (with 12 “undifferentiated”) (FDA, 2015).

Many labs have developed specific tests that they must validate and perform in house. These laboratory-developed tests (LDTs) are regulated by the Centers for Medicare and Medicaid (CMS) as high-complexity tests under the Clinical Laboratory Improvement Amendments of 1988 (CLIA '88). LDTs are not approved or cleared by the U. S. Food and Drug Administration; however, FDA clearance or approval is not currently required for clinical use.

***\*\*\*Note: This Medical Policy is complex and technical. For questions concerning the technical language and/or specific clinical indications for its use, please consult your physician.***

## Policy

---

**BCBSNC will provide coverage for Human Immunodeficiency Virus testing when it is determined the medical criteria or reimbursement guidelines below are met.**

## Benefits Application

---

This medical policy relates only to the services or supplies described herein. Please refer to the Member's Benefit Booklet for availability of benefits. Member's benefits may vary according to benefit design; therefore member benefit language should be reviewed before applying the terms of this medical policy.

## When Human Immunodeficiency Virus testing is covered

---

Reimbursement for initial screening for HIV infection for individuals 11 to 65 years of age is allowed.

Reimbursement for repeat screening for HIV infection (no less than 90 days after initial screening) for individuals 11 to 65 years of age is allowed.

Reimbursement for HIV genotyping or phenotyping is allowed for **any** of the following situations:

- a) Prior to initiating doravirine therapy (genotyping and phenotyping is **required**).
- b) For individuals who have failed a course of antiviral therapy.
- c) For individuals who have suboptimal viral load reduction.
- d) For individuals who have been noncompliant with therapy.
- e) To guide treatment decisions in individuals with acute or recent infection (within the last 6 months).
- f) For antiretroviral naïve individuals entering treatment.

# Human Immunodeficiency Virus – M2116

- g) For all HIV-infected pregnant individuals in the following situations:
  - i. Before initiation of antiretroviral therapy
  - ii. For those with detectable HIV RNA levels

Reimbursement for HIV phenotyping is allowed for treatment-experienced individuals on failing regimens who are thought to have multidrug resistance.

Reimbursement is allowed for a baseline HIV quantification when the risk of HIV infection is significant and the initiation of therapy is expected in **any** of the following situations:

- a) In an at-risk individual with persistence of borderline or equivocal serologic reactivity.
- b) In an at-risk individual with signs and symptoms of acute retroviral syndrome (characterized by fever, malaise, lymphadenopathy, and rash).

Reimbursement is allowed for plasma quantification of HIV-1 RNA or HIV-2 RNA (see Note 1) for **any** of the following situations:

- a) For monitoring disease progression in HIV-infected individuals.
- b) For monitoring response to antiretroviral therapy.
- c) For infants younger than 18 months born to HIV-positive mothers (antibody tests may be confounded by maternal antibodies in this time frame).
- d) For predicting maternal-fetal transmission of HIV-1 or HIV-2.

## When Human Immunodeficiency Virus testing is not covered

---

Reimbursement is not allowed for routine use of combined genotyping and phenotyping.

Reimbursement is not allowed for drug susceptibility phenotype prediction using genotypic comparison to known genotypic/phenotypic database.

---

### NOTES:

**Note 1:** Because differences in absolute HIV copy number are known to occur using different assays, plasma HIV RNA levels should be measured by the same analytical method. A change in assay method may necessitate re-establishment of a baseline.

## Table of Terminology

Term	Definition
AAP	American Academy of Pediatrics
AI/A1	Strong panel support – Evidence from $\geq 1$ RCTs published in the peer-reviewed literature or presented in abstract form at peer-reviewed scientific meetings
ACOG	American College of Obstetricians and Gynecologists
AIDS	Acquired Immune Deficiency Syndrome
AII/A2	Strong panel support - Data from well-designed nonrandomized trials or observational cohort studies with long-term clinical outcomes
AIIa	Strong panel support – Evidence from cohort or case-control studies published in the peer-reviewed literature
AIII	Strong panel support – Based on the panel’s analysis of the available evidence
ART	Antiretroviral treatment (also refers in some instances to antiretroviral testing and antiretroviral therapy)



# Human Immunodeficiency Virus – M2116

ARV	Antiretroviral
ASHM	The Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine
ATCC	American Type Culture Collection
BHIVA	British HIV Association
BII/B2	Moderate panel support - Data from well-designed nonrandomized trials or observational cohort studies with long-term clinical outcomes
BIIa	Moderate panel support – Evidence from cohort or case-control studies published in the peer-reviewed literature
BIII	Moderate panel support – Based on the panel’s analysis of the available evidence
CCR5	C-C chemokine receptor type 5
CD4	Cluster of differentiation 4
CDC	Centers for Disease Control and Prevention
CIII	Limited or weak panel support – Based on the panel’s analysis of the available evidence
CMS	Centers for Medicare and Medicaid
CPD	Citrate-phosphate-dextrose
CSF	Cerebrospinal fluid
CTM	COBAS TaqMan
DHHS	Department Of Health and Human Services
DNA	Deoxyribose nucleic acid
EACS	European Acquired Immune Deficiency Syndrome Clinical Society
EDTA	Ethylenediaminetetraacetic acid
FDA	Food and Drug Administration
GIS	Genotypic interpretation systems
GPP	General practice point
GT	Genotype
HIV	Human Immunodeficiency Virus
HIV-1	Human Immunodeficiency Virus, Type 1
HIV-2	Human Immunodeficiency Virus, Type 2
HIVDR	HIV drug resistance
HIVMA	HIV Medicine Association
HIV-VL	HIV viral load
IDSA	Infection Diseases Society of America
INSTI	Integrase strand transfer inhibitor
K103N	Lysine to aspartate polymorphism
LADR V	Low abundant drug resistant variant
LDT	Laboratory developed test
NAAT	Nucleic acid amplification test
NGS	Next generation sequencing
NNRTIs	Non-nucleoside reverse transcriptase inhibitors
NRTIs	Nucleoside reverse transcriptase inhibitors
NYSDOH	New York State Department of Health
PCR	Polymerase chain reaction
PEP	Postexposure prophylaxis
PIs	Protease inhibitors
PR	Protease
RAL	Raltegravir
RCT	Randomized controlled trial

# Human Immunodeficiency Virus – M2116

RNA	Ribonucleic acid
RT	Reverse transcriptase
RT-PCR	Reverse transcriptase polymerase chain reaction
RVA	Recombinant virus assay
SMFM	Society for Maternal-Fetal Medicine
SS	Sanger sequencing
TDR	Total drug resistance
USPSTF	United States Preventive Services Task Force

## Policy Guidelines

### Guidelines and Recommendations

#### Department of Health and Human Services(DHHS )

The Department of Health and Human Services (DHHS) (DHHS) Panel on Antiretroviral Guidelines for Adults and Adolescents updated the guidelines on use of antiretroviral drugs in 2022. The panel states “viral load is the most important indicator of initial and sustained response to ART and should be measured in all HIV-infected patients at entry into care (AI), at initiation of therapy (AI), and on a regular basis thereafter. Pre-treatment viral load level is also an important factor in the selection of an initial ARV regimen because several currently approved ARV drugs or regimens have been associated with poorer responses in patients with high baseline viral load.”

The panel’s recommendations on the frequency of viral load monitoring are summarized below (DHHS, 2022a):

- “After initiation of ART: Plasma viral load should be measured before initiation of ART and within 4 to 8 weeks after treatment initiation (AIII). The purpose of the measurements is to confirm an adequate virologic response to ART, indicating appropriate regimen selection and patient adherence to therapy. Repeat viral load measurement should be performed at 4- to 8-week intervals until the level falls below the assay’s limit of detection (BIII).”
- “In patients with viral suppression, with ART modification because of drug toxicity or for regimen simplification: Viral load measurement should be performed within 4 to 8 weeks after changing therapy (AIII). The purpose of viral load monitoring at this point is to confirm the effectiveness of the new regimen.”
- “In patients on a stable, suppressive ARV regimen: Viral load measurement should be repeated every 3 to 4 months (AIII) or as clinically indicated to confirm continuous viral suppression. Clinicians may extend the interval to 6 months for adherent patients whose viral load has been suppressed for more than a year, whose clinical and immunologic status is stable, and who are not at risk for inadequate adherence (AIII).”
- “In patients with virologic failure who require a change in ARV regimen: Plasma viral load should be measured before ART change and within 4 to 8 weeks after treatment modification (AIII). The purpose of the measurements is to confirm an adequate virologic response to the new regimen. Repeat viral load measurement should be performed at 4- to 8-week intervals until the level falls below the assay’s limit of detection (BIII). If viral suppression is not possible, repeat viral load measurement every 3 months or more frequently if indicated (AIII).”
- “In patients with suboptimal response: The frequency of viral load monitoring will depend on clinical circumstances, such as adherence and availability of further treatment options. In addition to viral load monitoring, several other factors—such as patient adherence to prescribed medications, suboptimal drug exposure, or drug interactions—should be assessed. Patients who fail to achieve viral suppression should undergo drug-resistance testing to aid in the selection of an alternative ARV regimen.”

# Human Immunodeficiency Virus – M2116

The guideline also comments on HIV-2. Although the optimal treatment strategy has not been defined, the guideline does recommend that quantitative plasma HIV-2 RNA viral load testing should be performed before initiating ART (AIII). HIV-2 RNA should also be used to assess treatment response. The guideline also notes that the “Genius HIV 1/2 Supplemental Assay (Bio-Rad Laboratories)” is FDA-approved to differentiate HIV-1 infection from HIV-2 infection (DHHS, 2022a).

The CDC refers to the above guidelines on their website (CDC, 2021).

In an updated review in 2022, the DHHS also strongly recommended (AIII) that “A blood sample for genotypic testing should be sent to the laboratory before initiation of ART.” Moreover, “Pregnancy testing should be performed in persons of childbearing potential before initiation of ART.”

The DHHS propounds further, stating the following:

- “Combination immunoassays that detect HIV-1 and HIV-2 antibodies and HIV p24 antigen (Ag/Ab assays) are part of the recommended initial laboratory HIV testing algorithm, primarily due to their enhanced ability to detect acute HIV infection. Specimens that are reactive on an initial Ag/Ab assay should be tested with an immunoassay that differentiates HIV-1 from HIV-2 antibodies. Specimens that are reactive on the initial assay and have either negative or indeterminate antibody differentiation test result should be tested for quantitative or qualitative HIV RNA; an undetectable HIV RNA test result indicates that the original Ag/Ab test result was a false positive. Detection of HIV RNA in this setting indicates that acute HIV infection is highly likely.”
- “HIV infection should be confirmed by repeat quantitative HIV RNA testing or subsequent testing to document HIV antibody seroconversion.”
- “The proposed threshold of <3,000 copies/mL is based on historical data that used laboratory methods that are now considered obsolete.<sup>18</sup> These older viral load assays demonstrated false-positive cases of acute HIV infection at HIV RNA levels of <3,000 copies/mL. However, improvements in plasma viral load methodology suggest that any positive result on a quantitative plasma HIV RNA test in the setting of a negative or indeterminate antibody test result is highly consistent with acute HIV infection, including at HIV RNA levels of <3,000 copies/mL. HIV RNA levels in acute infection are generally very high (e.g., >100,000 copies/mL);<sup>1, 2, 4</sup> however, levels may be <3,000 copies/mL in the earliest weeks following infection as viral load continues to rise. Therefore, when a low-positive quantitative HIV RNA test result is present at this level, the HIV RNA test should be repeated on a new blood specimen to confirm the diagnosis. Repeated false-positive HIV RNA test results are unlikely” (DHHS, 2022a).

As persons who acquire HIV while taking pre-exposure prophylaxis (PrEP) may present ambiguous HIV test results, the DHHS proposes that:

- “A positive HIV Ag/Ab test result or a positive HIV RNA test result in the setting of a negative HIV antibody test result should prompt immediate confirmation of HIV diagnosis. It is important to collect a new blood specimen to verify the HIV diagnosis before initiating ART.”
- “In people with HIV RNA level  $\geq 200$  copies/mL who are taking PrEP, immediate initiation of an effective HIV treatment regimen is recommended while awaiting confirmation of HIV diagnosis (AIII).”
- “In people taking PrEP who have a negative HIV antibody test result and a very low-positive quantitative HIV RNA test result ( $\geq 200$  copies/mL) a confirmatory HIV antibody test and repeat quantitative plasma HIV RNA test should be performed, and results should be available before initiating ART.”

# Human Immunodeficiency Virus – M2116

- “In rare cases, particularly when PrEP is transitioned to an antiretroviral (ARV) regimen and HIV RNA and antibody diagnostic testing are inconclusive, HIV DNA testing may be of value” (DHHS, 2022a).

The Department of Health and Human Services (DHHS, 2022a, 2022b, 2022c) updated their guidelines for using drug resistance assays in HIV infections. The guidelines recommend HIV genotyping or phenotyping in the following situations among pregnant individuals and reducing perinatal HIV transmission in the US:

- “General Principles Regarding Use of Antiretroviral Drugs During Pregnancy
  - Antiretroviral (ARV) drug-resistance genotype evaluations or assays should be performed before starting ARV drug regimens in people who are ARV-naive (AII) or ARV-experienced (AIII) and before modifying ARV drug regimens (AII) in people whose HIV RNA levels are above the threshold for resistance testing (i.e., >500 copies/mL to 1,000 copies/mL).
  - In pregnant people who are not already receiving ART, ART should be initiated before results of drug resistance testing are available because earlier viral suppression has been associated with lower risk of transmission. When ART is initiated before results are available, the regimen should be modified, if necessary, based on resistance assay results (AII).”
- “Pregnant People with HIV Who Have Never Received Antiretroviral Drugs (Antiretroviral Naive)
  - The results of ARV drug-resistance studies should guide the selection of ARV regimens in people whose HIV RNA levels are above the threshold for resistance testing (i.e., >500 copies/mL to 1,000 copies/mL) (see Antiretroviral Drug Resistance and Resistance Testing in Pregnancy) (AII). However, ART initiation should not be delayed while awaiting results of resistance testing. When ART is initiated before the results of the drug resistance assays are available, the ARV regimen should be modified, if necessary, based on the resistance assay results (AII).”
- “Pregnant People with HIV Who Are Currently Receiving Antiretroviral Therapy
  - ARV drug-resistance testing should be performed to assist the selection of active drugs when changing ARV regimens in pregnant people who are experiencing virologic failure on ART and who have HIV RNA levels >500 copies/mL to 1,000 copies/mL (AII). In individuals who have HIV RNA levels >500 copies/mL but <1,000 copies/mL, testing may be unsuccessful but still should be considered (BII).”
- “Pregnant People with HIV Who Have Previously Received Antiretroviral Medications but Are Not Currently Receiving Any Antiretroviral Medications
  - If HIV RNA is above the threshold for standard genotypic drug resistance testing (i.e., >500 to 1,000 copies/mL), ARV drug-resistance testing should be performed prior to starting an ARV drug regimen (AIII)
  - ART should be initiated prior to receiving results of current ARV resistance assays. ART should be modified based on the results of the resistance assay, if necessary (AII).”
- “Monitoring during Pregnancy
  - HIV drug-resistance testing (genotypic testing and, if indicated, phenotypic testing) should be performed during pregnancy in those whose HIV RNA levels are above the threshold for resistance testing (i.e., >500 copies/mL to 1,000 copies/mL) before –
    - Initiating ART in antiretroviral (ARV)-naive pregnant people who have not been previously tested for ARV drug resistance (AII);
    - Initiating ART in ARV-experienced pregnant people (including those who have received preexposure prophylaxis) (AIII); or

# Human Immunodeficiency Virus – M2116

- Modifying ARV regimens for people with HIV who become pregnant while receiving ARV drugs or people who have suboptimal virologic response to ARV drugs that were started during pregnancy (AII).
- ART should be initiated in pregnant patients prior to receiving the results of ARV-resistance tests. ART should be modified, if necessary, based on the results of resistance testing (AII).”
- “Antiretroviral Drug Resistance and Resistance Testing in Pregnancy
  - HIV drug-resistance testing (genotypic and, if indicated, phenotypic) should be performed in persons living with HIV whose HIV RNA levels are above the threshold for resistance testing (i.e., >500 to 1,000 copies/mL) before
    - Initiating ART in ARV-naive pregnant women who have not been previously tested for ARV resistance (AII),
    - initiating ART in ARV-experienced pregnant women (including those who have received pre-exposure prophylaxis) (AIII), or
    - modifying ART regimens for those who are newly pregnant and receiving ARV drugs or who have suboptimal virologic response to the ARV drugs during pregnancy (AII).
  - Phenotypic resistance testing is indicated for treatment-experienced persons on failing regimens who are thought to have multidrug resistance (BIII).
  - ART should be initiated in pregnant persons before receiving results of ARV-resistance testing; ART should be modified, if necessary, based on the results of resistance assays (AII).
  - If the use of an integrase strand transfer inhibitor (INSTI) is being considered and INSTI resistance is a concern, providers should supplement standard resistance testing with a specific INSTI genotypic resistance assay (AIII). INSTI resistance may be a concern if-
    - a patient received prior treatment that included an INSTI, *or*
    - a patient has a history with a sexual partner on INSTI therapy who was not virologically suppressed or with unknown viral load
    - documented zidovudine (ZDV) resistance does not affect the indications for use of intrapartum intravenous ZDV (BIII).” (DHHS, 2022c).

Among adults and adolescents living with HIV, the DHHS recommends the following for drug resistance testing:

- “For initial treatment:
  - HIV drug-resistance testing is recommended at entry into care for persons with HIV to guide selection of the initial antiretroviral therapy (ART) regimen (AII). If therapy is deferred, repeat testing may be considered at the time of ART initiation (CIII)
  - Genotypic, rather than phenotypic, testing is the preferred resistance testing to guide therapy in antiretroviral (ARV)-naive patients (AIII)
  - In persons with acute or recent (early) HIV infection, in pregnant people with HIV, or in people who will initiate ART on the day of or soon after HIV diagnosis, ART initiation should not be delayed while awaiting resistance testing results; the regimen can be modified once results are reported (AIII)
  - Standard genotypic drug-resistance testing in ARV-naive persons involves testing for mutations in the reverse transcriptase and protease genes. If transmitted integrase strand transfer inhibitor (INSTI) resistance is suspected or if the person has used long-acting cabotegravir (CAB-LA) as pre-exposure prophylaxis (PrEP) in the past, providers should ensure that genotypic resistance testing also includes the integrase gene (AIII).
- For Antiretroviral Therapy-Experienced Persons:

# Human Immunodeficiency Virus – M2116

- HIV drug-resistance testing should be performed to assist the selection of active drugs when changing ART regimens in the following patients:
  - People with virologic failure and HIV-RNA levels >200 copies/mL (AI for >1,000 copies/mL, AIII for 501–1,000 copies/mL, CIII for confirmed HIV RNA 201–500 copies/mL). For people with confirmed HIV-RNA levels >200 copies/mL but <500 copies/mL, drug-resistance testing may be unsuccessful but should still be considered.
  - Persons with suboptimal viral load reduction (AII)
- Reverse transcriptase and protease genotypic resistance testing should be performed on everyone with virologic failure; integrase resistance testing (which may need to be ordered separately) should be performed on individuals experiencing virologic failure while receiving an INSTI-based regimen (AII).
- For persons taking a non–long-acting ARV regimen, drug-resistance testing in the setting of virologic failure should be performed while the person is still taking their ARV regimen or, if that is not possible, within 4 weeks after discontinuing their ARV regimen (AII). If more than 4 weeks have elapsed since the non–long-acting agents were discontinued, resistance testing may still provide useful information to guide therapy; however, it is important to recognize that previously-selected resistance mutations can be missed due to lack of drug-selective pressure (CIII).
- Given the long half-lives of the long-acting injectable ARV drugs, resistance testing (including testing for resistance to INSTIs) should be performed in all persons who have experienced virologic failure on a regimen of long-acting CAB and rilpivirine or acquired HIV after receiving CAB-LA as PrEP, regardless of the amount of time since drug discontinuation (AIII).
- Genotypic testing is preferred over phenotypic resistance testing to guide therapy in people with suboptimal virologic response or virologic failure while on first- or second-line regimens and in people in whom resistance mutation patterns are known or not expected to be complex (AII).
- The addition of phenotypic- to genotypic resistance testing is recommended for people with known or suspected complex drug resistance mutation patterns (BIII).
- All prior and current drug-resistance test results, if available, should be reviewed and considered when constructing a new regimen for a patient (AIII)” (DHHS, 2022a).

In terms of the usage of drug-resistance assays among adolescents and adults with HIV, the DHHS recommends the following:

- “In acute or recent (early) HIV infection: Drug-resistance testing is recommended (AII). A genotypic assay is generally preferred (AIII). Treatment should not be delayed while awaiting results of resistance testing (AIII).
  - If ART is deferred, repeat resistance testing may be considered when therapy is initiated (CIII). A genotypic assay is generally preferred (AIII).”
- “In ART-naïve patients with chronic HIV: Drug-resistance testing is recommended at entry into HIV care to guide selection of initial ART (AII). A genotypic assay is generally preferred.”
  - For pregnant persons, or if ART will be initiated on the day of or soon after HIV diagnosis, treatment can be initiated prior to receiving resistance testing results.
  - If an INSTI is considered for an ART-naïve patient and/or transmitted INSTI resistance is a concern, providers should supplement standard resistance testing with a specific INSTI genotypic resistance assay, which may need to be ordered separately (AIII).

# Human Immunodeficiency Virus – M2116

- If therapy is deferred, repeat resistance testing may be considered when therapy is initiated (CIII). A genotypic assay is generally preferred (AIII).”
- “In patients with virologic failure: Drug-resistance testing is recommended in patients on combination ART with HIV RNA levels >200 copies/mL (AI for >1,000 copies/mL, AIII for 501–1,000 copies/mL) and a confirmed HIV RNA 201–500 copies/mL (CIII). In patients with confirmed HIV RNA levels between 201 - 500 copies/mL, testing may not be successful but should still be considered.
  - “Resistance testing should be done while the patient is taking ART or, if that is not possible, within 4 weeks after ART discontinuation of non – long-acting ARV drugs (AII). If >4 weeks have elapsed, resistance testing may still be useful to guide therapy; however, previously-selected mutations can be missed due to lack of drug selective pressure (CIII).”
  - “A standard genotypic resistance assay is generally preferred for patients experiencing virologic failure on their first or second ARV regimens and for those with expected noncomplex resistance patterns (AII).”
  - “All prior and current drug-resistance testing results should be reviewed and considered when designing a new ARV.”
  - “When virologic failure occurs in a patient on an INSTI based regimen or in a patient with a history of INSTI use, genotypic testing for INSTI resistance should be performed to determine whether to include drugs from this class in subsequent regimens (AII).”
  - “Adding phenotypic testing to genotypic testing is generally preferred in patients with known or suspected complex drug-resistance patterns (BIII)”
- “In patients with suboptimal suppression of viral load: Drug resistance testing is recommended in patients with suboptimal viral load suppression after initiation of ART. (AII)”
- “In pregnant people with HIV: Genotypic resistance testing is recommended for all pregnant people before initiation of ART (AIII) and for those entering pregnancy with detectable HIV RNA levels while on therapy. (AI)”
- “In patients with Undetectable Viral Load or Low-Level Viremia Who Are Planning to Change Their ARV Regimen HIV-1: proviral DNA resistance assays may be useful in patients with HIV RNA below the limit of detection or with low-level viremia, where a HIV RNA genotypic assay is unlikely to be successful (CIII)” (DHHS, 2022a)

The DHHS also added guidelines on genotypic and phenotypic testing for pediatric HIV infection:

- “Antiretroviral (ARV) drug- resistance testing is recommended at the time of HIV diagnosis, before initiation of therapy, in all ART naive patients, and before switching regimens in patients with treatment failure (AII). Genotypic resistance testing is preferred for this purpose (AIII).”
- “Phenotypic resistance testing should be considered (usually in addition to genotypic resistance testing) for patients with known or suspected complex drug resistance mutation patterns, which generally arise after a patient has experienced virologic failure on multiple ARV regimens (CIII)” (DHHS, 2022b).

## International Antiviral Society

The International Antiviral Society published a 2022 update titled “Antiretroviral Drugs for Treatment and Prevention of HIV Infection in Adults.” The guideline also recommends laboratory testing to “characterize” the HIV stage prior to starting antiretroviral testing (ART); this is done by assessing HIV RNA level (Gandhi et al., 2022).

# Human Immunodeficiency Virus – M2116

The guideline also remarks on the frequency of testing during ART. Their recommendations are as follows:

- “Within 6 weeks of starting ART, assessment of treatment adherence and tolerability is recommended, along with the measurement of HIV RNA level.”
- “If the HIV RNA level has not declined by 2 log<sub>10</sub> copies/mL within 12 weeks of therapy and adherence appears to be sufficient, then a genotype based on the patient’s regimen is recommended.”
- “If the patient remains virally suppressed, clinically stable, and adherent to medications, then HIV RNA levels should be monitored every 3 months until virally suppressed for at least 1 year. Afterward, the frequency of viral monitoring can be changed to every 6 months.”
- “If HIV RNA level is greater than 200 copies/mL on 2 consecutive measurements, then HIV RT-pro genotype and InSTI [in integrase strand transfer inhibitor]genotype (if the patient was receiving an InSTI) testing are recommended.”
- “For patients with intermittent or persistent low-level viremia between 50 and 200 copies/mL, assessments for ART adherence, tolerability, and toxic effects are recommended, but changing ART regimens is not recommended unless ART toxicity or intolerance are identified” (Gandhi et al., 2022).

On resistance test, the 2022 update notes that, “in persons diagnosed with HIV while receiving TXF-based PrEP, resistance testing should be performed but initiation of ART need not be delayed while awaiting genotype results.” The panel further recommends:

- “Unless there is documented or suspected history of treatment failure, proviral resistance testing is not required prior to switching to 2-drug therapy, even if there is no available pretreatment resistance test result.”
- “For patients who have maintained viral suppression, switching from long-acting injectable cabotegravir plus rilpivirine back to daily oral therapy can be done without the need for proviral DNA resistance testing.”
- “If virologic failure is confirmed, genotype resistance testing should be performed, preferably while patients are taking the failing therapy. Resistance testing is still recommended even if a regimen has been discontinued or a person acknowledges poor medication adherence” (Gandhi et al., 2022).

## **Infectious Diseases Society of America (IDSA)**

The IDSA recommends that “A quantitative HIV RNA (viral load) level should be obtained upon initiation of care (strong recommendation, high quality evidence)” (Thompson et al., 2020).

IDSA recommends rechecking HIV RNA after 2-4 weeks of initiating antiretroviral therapy (ART) (and no later than 8 weeks). From there, IDSA recommends “checking HIV RNA every 4-8 weeks until suppression is achieved.” The IDSA also notes that viral load “should” be monitored every 3-4 months to “confirm maintenance of suppression below the limit of assay detection,” 6 months for “adherent patients whose viral load has been suppressed for more than 2 years and whose clinical and immunologic status is stable”, and more frequently after initiation or change in ART (IDSA recommends within 2-4 weeks of initiation or change but not more than 8 weeks) (Thompson et al., 2020).

Overall, IDSA lists two primary uses for viral load testing; to establish baseline and to monitor viral suppression (Thompson et al., 2020).

## **American College of Obstetricians and Gynecologists (ACOG)**



# Human Immunodeficiency Virus – M2116

In 2014, ACOG released “Committee on Gynecologic Practice: Routine human immunodeficiency virus screening,” which they reaffirmed in 2020. Regarding routine human immunodeficiency screening, “The American College of Obstetricians and Gynecologists (the College) recommends routine HIV screening for females aged 13–64 years and older women with risk factors. Screening after age 64 years is indicated if there is ongoing risk of HIV infection, as indicated by risk assessment (eg, new sexual partners)” (ACOG, 2014).

The College also expatiates upon repeat testing, entrusting obstetrician–gynecologists to annually review patients’ risk factors for HIV and assess their needs, and recommends that “HIV testing should be offered at least annually to women who

- are injection drug users
- are sex partners of injection-drug users
- exchange sex for money or drugs
- are sex partners of HIV-infected persons
- have had sex with men who have sex with men since the most recent HIV test
- have had more than one sex partner since their most recent HIV test

The opportunity for repeat testing should be made available to all women even in the absence of identified risk factors. Repeat screening after age 64 years is indicated if there is ongoing risk of HIV infection, as indicated by an individualized risk assessment. Obstetrician–gynecologists also should encourage women and their prospective sex partners to be tested before initiating a new sexual relationship. The benefits of periodic retesting should be discussed with patients and provided if requested, regardless of risk factors. Patients may be concerned about their status and do not know about or want to disclose risk-taking behavior to their health care providers” (ACOG, 2014).

In their 2018 committee opinion “Labor and Delivery Management of Women With Human Immunodeficiency Virus Infection,” ACOG notes that current and ongoing research has shown that “treatment of HIV-infected pregnant women with combined antiretroviral therapy can achieve a 1–2% or lower risk of mother-to-child transmission if maternal viral loads of 1,000 copies/mL or less can be sustained, independent of the route of delivery or duration of ruptured membranes before delivery.” ACOG further observes that “the risk of mother-to-child transmission in HIV-infected women with high viral loads can be reduced by performing cesarean deliveries before the onset of labor and before rupture of membranes [cesarean delivery in this document [the ACOG guideline]], in conjunction with the use of peripartum maternal antiretroviral therapy.”

ACOG recommends offering a “scheduled prelabor cesarean delivery at 38 0/7 weeks of gestation to reduce the risk of mother-to-child transmission” if an HIV-positive pregnant woman is found to have a viral load of over 1000 copies/mL at or near delivery, independent of antepartum antiretroviral therapy. This recommendation also applies to patients whose viral load is unknown (ACOG, 2018).

## **Society for Maternal-Fetal Medicine (SMFM)**

The SMFM published a “checklist for pregnancy management in persons with HIV.” Although these checklists are not definitive, they are intended to “help ensure that all relevant elements are considered for every person with HIV during prepregnancy, antepartum, intrapartum, and postpartum periods.” During the third trimester, the checklist calls for viral load to be assessed at 34–36 weeks for delivery planning (and to assess adherence and viral resistance if viral load is not suppressed). Further, if the viral load is found to be  $\geq 1000$  copies/mL at 37–38 weeks, a cesarean delivery should be scheduled for 38 weeks (Gibson & Toner, 2020).

# Human Immunodeficiency Virus – M2116

## British HIV Association

The British HIV Association (BHIVA) makes several recommendations regarding assessment of viral load during the routine investigation and/or maintenance of HIV-1 positive adults. Relevant recommendations are as follows:

- “We recommend that an HIV viral load should be performed at the first visit following serological diagnosis (1A).
- We recommend that undetectable viral load result whilst not on treatment needs repeating, review of serology to exclude HIV-2 and measurement on a different viral load assay (1D).
- We recommend a repeat HIV viral load in all new transfers prior to repeat prescriptions if it is not possible to confirm a recent viral load from the previous clinic (1A).
- We recommend that viral load measurements be taken at 1, 3 and 6 months after starting ART (1B).
- We recommend that additional viral load measurements are taken between 2 and 5 months after starting ART if viral load has not decreased at least 10-fold after 1 month of ART or there are concerns about the patient’s adherence to therapy (1D).
- We recommend that viral load testing should be performed routinely every 6 months (1A) and might be at intervals of up to 12 months for patients established on ART that includes a PI (GPP) [general practice point].
- We recommend that viral load rebound to above 50 copies/mL should be confirmed by testing a subsequent sample (2A). Repeat testing of the same sample is not recommended.
- For patients stable on ART we recommend that:
  - Frequent (3–4 monthly) viral load follow-ups of individuals with stable unsuppressed (<200 copies/mL) viral loads if they are managed as low-level viraemic patients according to the BHIVA treatment guidelines (1D).
  - CSF HIV viral load measurement should be considered to exclude compartmentalisation (1C)” (BHIVA, 2019).

The BHIVA released guidelines for the management of HIV-2 (BHIVA, 2021). For the diagnosis of HIV-2, the BHIV recommends:

For the diagnosis of chronic HIV-2:

- “An initial diagnosis of chronic HIV-2 infection should be made using a total of three CE-marked serology tests (i.e. tests conform to EU health and safety requirements) performed in an ISO 15189-accredited laboratory. There must be reactivity in two CE-marked fourth-generation tests for HIV-1 and HIV-2, followed by differentiation of HIV-2 by a third CE-marked antibody-only test.”
- “Clinicians should consider revisiting a previous diagnosis of HIV-1 by repeating HIV-2 serology and molecular tests in individuals with an undetectable HIV-1 viral load in the absence of ART, but a falling CD4 count. This is in order to detect the possibility of missed HIV-1 and HIV-2 dual infection.”
- “In those with diagnosed HIV-2 with an undetectable viral load in the absence of ART, clinicians should consider repeating HIV-1 diagnostic tests, if their CD4 count falls. This is to investigate the possibility of HIV-1 superinfection.”

For the diagnosis of acute primary HIV-2

- “Investigation for acute or very recent HIV-2 infection should start as for diagnosis of chronic HIV-2 infection. A negative HIV-2 screening result on a blood sample taken within 3 months of the likely exposure should be further investigated at 6 weeks and 3 months, with parallel testing for HIV-2 viral RNA and, if necessary, HIV-2 proviral DNA.”

# Human Immunodeficiency Virus – M2116

For the investigation of indeterminate HIV-1 or HIV-2:

- “We recommend that any HIV-1 or HIV-2 serology that does not fit into a clear pattern of a confirmed laboratory diagnosis is fully investigated for the presence or absence of HIV-2 infection, and that this should be established by PCR for HIV-2 proviral DNA.”

For measuring HIV-2 viral load:

- “If the pre-treatment viral load was detectable, the viral load should be measured at 1, 3 and 6 months after starting or changing ART and then 3–6 monthly.
- “If the pre-treatment viral load was undetectable, the viral load should be measured at 1 month and then 6 monthly.”
- The HIV-2 viral load should be repeated in those on ART when it has been maximally suppressed and then becomes detectable.”
- Testing for drug resistance should be performed in those on ART when the HIV-2 viral load has been maximally suppressed and then becomes repeatedly detectable.”

For resistance testing:

- “Resistance testing should be performed at diagnosis, prior to treatment initiation and at virological failure, if the HIV-2 viral load meets the threshold of  $\geq 500$  copies/mL” (BHIVA, 2021).

## European AIDS Clinical Society (EACS)

The EACS recommends a genotypic resistance test to be ideally done at the time of HIV diagnosis; testing “should not delay ART initiation (it may be re-adjusted after genotypic test results). Resistance testing is also recommended to be performed in the setting of virological failure, “preferably on failing therapy (usually routinely available for HIV-VL levels  $>200$ - $500$  copies/mL and in specialized laboratories for lower levels of viremia) and obtain historical resistance testing for archived mutations.” For pregnant women, the EACS recommends performing resistance testing on women whose HIV-VL is not undetectable at third trimester, and “consider changing to or adding INSTI (RAL or DTG) if not on this class to obtain rapid HIV-VL decline.” When considering PEP, the EACS recommends resistance testing if the HIV-VL is detectable in an HIV-positive source person on ART. They also recommend baseline resistance testing when considering a combination regimen for ART-naïve children and adolescents living with HIV. Resistance testing should also be used to help guide the choice of treatment.

Additional genotypic recommendations include if the patient was not previously tested or if the patient is at risk of a super-infection. Genotypic resistance testing is also required prior to beginning treatment with doravirine. When switching strategies for “virologically suppressed persons,” Proviral DNA genotyping may be useful in persons with multiple virological failures, unavailable resistance history or low-level viremia at the time of switch. Results ought to be taken cautiously as proviral DNA genotype may not detect previous resistance mutations and can also detect clinically irrelevant mutations. Therefore, routine proviral DNA genotyping is currently not recommended.” The EACS recommends a genotypic test over a phenotypic test as genotype tests are more available and more sensitive (EACS, 2022).

## The American Academy of Pediatrics (AAP)

The AAP recommends:

- “Routine HIV screening is recommended for all youth 15 years or older, at least once, in health care settings.”

# Human Immunodeficiency Virus – M2116

- “After initial screening, youth at increased risk, including sexually active youth, should be rescreened at least annually, potentially as frequently as every 3 to 6 months if at high risk (male youth reporting male sexual contact, active injection drug users, transgender youth; having sexual partners who are HIV-infected, of both genders, or injection drug users; exchanging sex for drugs or money; or those who have had a diagnosis of or request testing for other STIs).”
- “Youth who request HIV screening at any time should be tested, even in the absence of reported risk factors” (Hsu & Rakhmanina, 2022)

The Bright Futures/AAP Periodicity Schedule describes the screenings, assessments, physical examinations, procedures, and timing of anticipatory guidance recommended for each age-related visit. These guidelines provide the following recommendation for HIV screening:

- STI/HIV screening annually starting at 11 years old, with at least one HIV screening between 15 and 18 (AAP, 2022).

## **United States Preventive Services Task Force (USPSTF)**

The USPSTF recommends screening adolescents under 15 who are at increased risk, adolescents and adults aged 15 to 65 years, and younger adolescents and older adults who are at increased risk should.

The USPSTF also recommends screening all pregnant women for HIV, including those in labor who are untested and whose HIV status is unknown (USPSTF, 2019). The CDC recognizes and supports these guidelines (CDC, 2020).

## **Billing/Coding/Physician Documentation Information**

---

This policy may apply to the following codes. Inclusion of a code in this section does not guarantee that it will be reimbursed. For further information on reimbursement guidelines, please see Administrative Policies on the Blue Cross Blue Shield of North Carolina web site at [www.bcbsnc.com](http://www.bcbsnc.com). They are listed in the Category Search on the Medical Policy search page.

*Applicable codes: 86689, 86701, 86702, 86703, 87389, 87390, 87391, 87534, 87535, 87536, 87537, 87538, 87539, 87806, 87900, 87901, 87903, 87904, 87906, 0219U, G0432, G0433, G0435, G0475, S3645*

BCBSNC may request medical records for determination of medical necessity. When medical records are requested, letters of support and/or explanation are often useful, but are not sufficient documentation unless all specific information needed to make a medical necessity determination is included.

## **Scientific Background and Reference Sources**

---

AAP. (2022). Bright Futures/AAP Recommendations for Preventive Pediatric Health Care (Periodicity Schedule). <https://www.aap.org/periodicityschedule>

ACOG. (2014). ACOG Committee Opinion no 596: Committee on Gynecologic Practice: Routine human immunodeficiency virus screening. *Obstet Gynecol*, 123(5), 1137-1139. <https://doi.org/10.1097/01.AOG.0000446828.64137.50>

# Human Immunodeficiency Virus – M2116

ACOG. (2018). ACOG Committee Opinion No. 751: Labor and Delivery Management of Women With Human Immunodeficiency Virus Infection. *Obstet Gynecol*, 132(3), e131-e137. <https://doi.org/10.1097/aog.0000000000002820>

ATCC. (2014). ATCC Teams with CDC and Thermo Fisher Scientific on Public Health RT-PCR Assay. [https://www.labbulletin.com/articles/ATCC-Teams-CDC-Thermo-Fisher-Scientific-Public-Health-RT-PCR-Assay/categories/20130120\\_11](https://www.labbulletin.com/articles/ATCC-Teams-CDC-Thermo-Fisher-Scientific-Public-Health-RT-PCR-Assay/categories/20130120_11)

Avram, C. M., Greiner, K. S., Tilden, E., & Caughey, A. B. (2019). Point-of-care HIV viral load in pregnant women without prenatal care: a cost-effectiveness analysis. *Am J Obstet Gynecol*, 221(3), 265.e261-265.e269. <https://doi.org/10.1016/j.ajog.2019.06.021>

BHIVA. (2019). BHIVA guidelines for the routine investigation and monitoring of adult HIV-1-positive individuals (2019 interim update). Retrieved 12/3/20 from <https://www.bhiva.org/file/DqZbRxfzLYtLg/Monitoring-Guidelines.pdf>

BHIVA. (2021). British HIV Association guidelines for the management of HIV-2 2021. <https://www.bhiva.org/file/615ee3de98539/BHIVA-guidelines-for-the-management-of-HIV-2.pdf>

BusinessWire. (2020). Aptima HIV-1 Quant Dx Assay Receives Additional FDA Approval for Use as an Aid in the Diagnosis of HIV Infection. <https://www.businesswire.com/news/home/20201120005242/en/>

Caliendo, A. (2022, March 12). Techniques and interpretation of HIV-1 RNA quantitation. <https://www.uptodate.com/contents/techniques-and-interpretation-of-hiv-1-rna-quantitation>

CDC. (2014). Revised surveillance case definition for HIV infection--United States, 2014. *MMWR Recomm Rep*, 63(Rr-03), 1-10. <https://www.cdc.gov/mmwr/preview/mmwrhtml/rr6303a1.htm>

CDC. (2020). HIV Preventative Services Coverage. <https://www.cdc.gov/nchhstp/highqualitycare/preventiveservices/hivaids.html>

CDC. (2021, September 21, 2021). HIV Treatment and Care. Retrieved 2/11/2021 from <https://www.cdc.gov/hiv/clinicians/treatment/treatment-clinicians.html>

Coffin, J., & Swanstrom, R. (2013). HIV Pathogenesis: Dynamics and Genetics of Viral Populations and Infected Cells. In *Cold Spring Harb Perspect Med* (Vol. 3). <https://doi.org/10.1101/cshperspect.a012526>

DHHS. (2022a). Guidelines for the Use of Antiretroviral Agents in Adults and Adolescents with HIV. <https://clinicalinfo.hiv.gov/sites/default/files/guidelines/documents/adult-adolescent-arv/guidelines-adult-adolescent-arv.pdf>

DHHS. (2022b, April 11, 2022). Guidelines for the Use of Antiretroviral Agents in Pediatric HIV Infection. <https://clinicalinfo.hiv.gov/en/guidelines/pediatric-arv/whats-new-guidelines>

DHHS. (2022c, March 17, 2022). Recommendations for the Use of Antiretroviral Drugs in Pregnant Women with HIV Infection and Interventions to Reduce Perinatal HIV Transmission in the United States. <https://clinicalinfo.hiv.gov/en/guidelines/perinatal/whats-new-guidelines>

EACS. (2022). European AIDS Clinical Society Guidelines Version 11.1 October 2022. [https://www.eacsociety.org/media/guidelines-11.1\\_final\\_09-10.pdf](https://www.eacsociety.org/media/guidelines-11.1_final_09-10.pdf)

# Human Immunodeficiency Virus – M2116

Ehret, R., Harb, K., Breuer, S., & Obermeier, M. (2022). Performance assessment of the new Xpert® HIV-1 viral load XC assay for quantification of HIV-1 viral loads. *J Clin Virol*, 149, 105127. <https://doi.org/10.1016/j.jcv.2022.105127>

FDA. (2007a). Abbott RealTime HIV-1 <https://www.fda.gov/media/73278/download>

FDA. (2007b). COBAS® AmpliPrep/COBAS® TaqMan® HIV-1 Test, Summary of Safety and Effectiveness <https://www.fda.gov/media/73824/download>

FDA. (2015). BioPlex 2200 HIV Ag-Ab Assay. <https://www.fda.gov/media/92862/download>

FDA. (2019). Geenius™ HIV 1/2 Supplemental Assay. <https://www.fda.gov/media/130312/download>

FDA. (2020). Aptima® HIV-1 Quant Dx Assay. Retrieved 1/22/2021 from <https://www.fda.gov/media/102425/download>

Fogel, J. M., Bonsall, D., Cummings, V., Bowden, R., Golubchik, T., de Cesare, M., Wilson, E. A., Gamble, T., del Rio, C., Batey, D. S., Mayer, K. H., Farley, J. E., Hughes, J. P., Remien, R. H., Beyrer, C., Fraser, C., & Eshleman, S. H. (2020). Performance of a high-throughput next-generation sequencing method for analysis of HIV drug resistance and viral load. *Journal of Antimicrobial Chemotherapy*, 75(12), 3510-3516. <https://doi.org/10.1093/jac/dkaa352>

Fox, Z. V., Geretti, A. M., Kjaer, J., Dragsted, U. B., Phillips, A. N., Gerstoft, J., Staszewski, S., Clotet, B., von Wyl, V., & Lundgren, J. D. (2007). The ability of four genotypic interpretation systems to predict virological response to ritonavir-boosted protease inhibitors. *Aids*, 21(15), 2033-2042. <https://doi.org/10.1097/QAD.0b013e32825a69e4>

Gandhi, R. T., Bedimo, R., Hoy, J. F., Landovitz, R. J., Smith, D. M., Eaton, E. F., Lehmann, C., Springer, S. A., Sax, P. E., Thompson, M. A., Benson, C. A., Buchbinder, S. P., del Rio, C., Eron, J. J., Jr, Günthard, H. F., Molina, J.-M., Jacobsen, D. M., & Saag, M. S. (2022). Antiretroviral Drugs for Treatment and Prevention of HIV Infection in Adults: 2022 Recommendations of the International Antiviral Society–USA Panel. *JAMA*. <https://doi.org/10.1001/jama.2022.22246>

Gibson, K. S., & Toner, L. E. (2020). Society for Maternal-Fetal Medicine Special Statement: Updated checklists for pregnancy management in persons with HIV. *American Journal of Obstetrics & Gynecology*, 223(5), B6-B11. <https://doi.org/10.1016/j.ajog.2020.08.064>

Gottlieb, G. (2022a, December 5). Clinical manifestations and diagnosis of HIV-2 infection. <https://www.uptodate.com/contents/clinical-manifestations-and-diagnosis-of-hiv-2-infection>

Gottlieb, G. (2022b). Epidemiology, transmission, natural history, and pathogenesis of HIV-2 infection. UpToDate. Retrieved 2/11/2022 from <https://www.uptodate.com/contents/epidemiology-transmission-natural-history-and-pathogenesis-of-hiv-2-infection>

Hopkins, M., Hau, S., Tiernan, C., Papadimitropoulos, A., Chawla, A., Beloukas, A., & Geretti, A. M. (2015). Comparative performance of the new Aptima HIV-1 Quant Dx assay with three commercial PCR-based HIV-1 RNA quantitation assays. *Journal of Clinical Virology*, 69, 56-62. <https://doi.org/https://doi.org/10.1016/j.jcv.2015.05.020>

# Human Immunodeficiency Virus – M2116

Hsu, K. K., & Rakhmanina, N. Y. (2022). Adolescents and Young Adults: The Pediatrician's Role in HIV Testing and Pre- and Postexposure HIV Prophylaxis. *Pediatrics*, 149(1). <https://doi.org/10.1542/peds.2021-055207>

Hughes, M. D., Johnson, V. A., Hirsch, M. S., Bremer, J. W., Elbeik, T., Erice, A., Kuritzkes, D. R., Scott, W. A., Spector, S. A., Basgoz, N., Fischl, M. A., & D'Aquila, R. T. (1997). Monitoring plasma HIV-1 RNA levels in addition to CD4+ lymphocyte count improves assessment of antiretroviral therapeutic response. ACTG 241 Protocol Virology Substudy Team. *Ann Intern Med*, 126(12), 929-938. <https://pubmed.ncbi.nlm.nih.gov/9182469/>

Kozal, M. (2019a, September 17). Interpretation of HIV drug resistance testing. <https://www.uptodate.com/contents/interpretation-of-hiv-drug-resistance-testing>

Kozal, M. (2019b, September 17). Overview of HIV drug resistance testing assays. <https://www.uptodate.com/contents/overview-of-hiv-drug-resistance-testing-assays>

LabCorp. (2021). Human Immunodeficiency Virus 1 (HIV-1) PhenoSense GT® Plus Integrase (Monogram® Phenotype + Genotype). <https://www.labcorp.com/tests/551920/human-immunodeficiency-virus-1-hiv-1-phenosense-gt-plus-integrase-monogram-phenotype-genotype>

Lindman, J., Hønge, B. L., Kjerulff, B., Medina, C., da Silva, Z. J., Erikstrup, C., Norrgren, H., & Månsson, F. (2019). Performance of Bio-Rad HIV-1/2 Confirmatory Assay in HIV-1, HIV-2 and HIV-1/2 dually reactive patients - comparison with INNO-LIA and immunocomb discriminatory assays. *J Virol Methods*, 268, 42-47. <https://doi.org/10.1016/j.jviromet.2019.03.005>

Mansky, L. M., & Temin, H. M. (1995). Lower in vivo mutation rate of human immunodeficiency virus type 1 than that predicted from the fidelity of purified reverse transcriptase. *J Virol*, 69(8), 5087-5094. <https://www.ncbi.nlm.nih.gov/pmc/articles/pmid/7541846/>

Mor, O., Gozlan, Y., Wax, M., Mileguir, F., Rakovsky, A., Noy, B., Mendelson, E., & Levy, I. (2015). Evaluation of the RealTime HIV-1, Xpert HIV-1, and Aptima HIV-1 Quant Dx Assays in Comparison to the NucliSens EasyQ HIV-1 v2.0 Assay for Quantification of HIV-1 Viral Load. *J Clin Microbiol*, 53(11), 3458-3465. <https://doi.org/10.1128/jcm.01806-15>

Pröll, J., Paar, C., Taylor, N., Skocic, M., Freystetter, A., Blaimschein, A., Mayr, R., Niklas, N., Atzmüller, S., Raml, E., & Wechselberger, C. (2022). New aspects of the Virus Life Cycle and Clinical Utility of Next Generation Sequencing based HIV-1 Resistance Testing in the Genomic, the Proviral and the Viral Reservoir of Peripheral Blood Mononuclear Cells. *Curr HIV Res*. <https://doi.org/10.2174/1570162x20666220324111418>

Quinn, T. (2022). Global epidemiology of HIV infection. UpToDate. <https://www.uptodate.com/contents/global-epidemiology-of-hiv-infection>

Raymond, S., Nicot, F., Abravanel, F., Minier, L., Carcenac, R., Lefebvre, C., Harter, A., Martin-Blondel, G., Delobel, P., & Izopet, J. (2020). Performance evaluation of the Vela Dx Sentosa next-generation sequencing system for HIV-1 DNA genotypic resistance. *Journal of Clinical Virology*, 122, 104229. <https://doi.org/https://doi.org/10.1016/j.jcv.2019.104229>

Rosemary, A., Chika, O., Jonathan, O., Godwin, I., Georgina, O., Azuka, O., Zaidat, M., Philippe, C., Oliver, E., Oche, A., David, O., Jay, S., Ibrahim, D., Mukhtar, A., Joshua, D., Chunfu, Y., Elliot, R., Beth, C., Phyllis, K., & Emmanuel, I. (2018). Genotyping

# Human Immunodeficiency Virus – M2116

performance evaluation of commercially available HIV-1 drug resistance test. *PLoS One*, 13(6), e0198246. <https://doi.org/10.1371/journal.pone.0198246>

Saag, M. S., Gandhi, R. T., Hoy, J. F., Landovitz, R. J., Thompson, M. A., Sax, P. E., Smith, D. M., Benson, C. A., Buchbinder, S. P., del Rio, C., Eron, J. J., Jr., Fätkenheuer, G., Günthard, H. F., Molina, J.-M., Jacobsen, D. M., & Volberding, P. A. (2020). Antiretroviral Drugs for Treatment and Prevention of HIV Infection in Adults: 2020 Recommendations of the International Antiviral Society–USA Panel. *JAMA*, 324(16), 1651-1669. <https://doi.org/10.1001/jama.2020.17025>

Saag, M. S., Holodniy, M., Kuritzkes, D. R., O'Brien, W. A., Coombs, R., Poscher, M. E., Jacobsen, D. M., Shaw, G. M., Richman, D. D., & Volberding, P. A. (1996). HIV viral load markers in clinical practice. *Nat Med*, 2(6), 625-629. <https://pubmed.ncbi.nlm.nih.gov/8640545/>

Sempa, J. B., Dushoff, J., Daniels, M. J., Castelnuovo, B., Kiragga, A. N., Nieuwoudt, M., & Bellan, S. E. (2016). Reevaluating Cumulative HIV-1 Viral Load as a Prognostic Predictor: Predicting Opportunistic Infection Incidence and Mortality in a Ugandan Cohort. *Am J Epidemiol*, 184(1), 67-77. <https://doi.org/10.1093/aje/kwv303>

Shen, C., Yu, X., Harrison, R. W., & Weber, I. T. (2016). Automated prediction of HIV drug resistance from genotype data. *BMC Bioinformatics*, 17 Suppl 8, 278. <https://doi.org/10.1186/s12859-016-1114-6>

Sollis, K. A., Smit, P. W., Fiscus, S., Ford, N., Vitoria, M., Essajee, S., Barnett, D., Cheng, B., Crowe, S. M., Denny, T., Landay, A., Stevens, W., Habiyambere, V., Perrins, J., & Peeling, R. W. (2014). Systematic review of the performance of HIV viral load technologies on plasma samples. *PLoS One*, 9(2), e85869. <https://doi.org/10.1371/journal.pone.0085869>

Swenson, L. C., Cobb, B., Geretti, A. M., Harrigan, P. R., Poljak, M., Seguin-Devaux, C., Verhofstede, C., Wirden, M., Amendola, A., Boni, J., Bourlet, T., Huder, J. B., Karasi, J. C., Zidovec Lepej, S., Lunar, M. M., Mukabayire, O., Schuurman, R., Tomazic, J., Van Laethem, K., . . . Wensing, A. M. (2014). Comparative performances of HIV-1 RNA load assays at low viral load levels: results of an international collaboration. *J Clin Microbiol*, 52(2), 517-523. <https://doi.org/10.1128/jcm.02461-13>

ThermoFisher. (2011). ViroSeq™ HIV-1 Genotyping System. [https://tools.thermofisher.com/content/sfs/manuals/cms\\_041134.pdf](https://tools.thermofisher.com/content/sfs/manuals/cms_041134.pdf)

Thompson, M. A., Horberg, M. A., Agwu, A. L., Colasanti, J. A., Jain, M. K., Short, W. R., Singh, T., & Aberg, J. A. (2020). Primary Care Guidance for Persons With Human Immunodeficiency Virus: 2020 Update by the HIV Medicine Association of the Infectious Diseases Society of America. *Clinical Infectious Diseases*. <https://doi.org/10.1093/cid/ciaa1391>

USPSTF. (2019). Screening for HIV Infection: US Preventive Services Task Force Recommendation Statement. Retrieved 23 from <https://doi.org/10.1001/jama.2019.6587>

Wood, B. R. (2021, March 8). The natural history and clinical features of HIV infection in adults and adolescents. UpToDate. <https://www.uptodate.com/contents/the-natural-history-and-clinical-features-of-hiv-infection-in-adults-and-adolescents>

Zhang, J., Rhee, S. Y., Taylor, J., & Shafer, R. W. (2005). Comparison of the precision and sensitivity of the Antivirogram and PhenoSense HIV drug susceptibility assays. *J Acquir Immune Defic Syndr*, 38(4), 439-444.



# Human Immunodeficiency Virus – M2116

## Policy Implementation/Update Information

---

- 1/1/19 New policy developed. BCBSNC will provide coverage for HIV genotyping and phenotyping when it is determined to be medically necessary because the medical criteria and guidelines are met. Medical Director review 1/1/2019. Policy noticed 1/1/2019 for effective date 4/1/2019. (sk)
- 9/10/19 Reviewed by Avalon 2nd Quarter 2019 CAB. Added “Related Policies” section. Policy Guidelines updated. References updated. Coding table removed from the Billing/Coding section of the policy. Medical Director review 8/2019. (sk)
- 10/29/19 Wording in the Policy, When Covered, and/or Not Covered section(s) changed from Medical Necessity to Reimbursement language, where needed. (gm)
- 3/10/20 Specialty Matched Consultant Advisory Panel review 2/19/2020. (sk)
- 7/28/20 Reviewed by Avalon 2nd Quarter 2020 CAB. Literature review updated. Not Covered section changed from Medical Necessity to Reimbursement language, where needed. Policy Guidelines updated. References updated. Medical Director review 7/2020. (bb)
- 3/9/21 Specialty Matched Consultant Advisory Panel review 2/17/2021. (sk)
- 8/24/21 Reviewed by Avalon 2nd Quarter 2021 CAB. Scientific Background updated. Policy Guidelines updated. New code 0219U added to Billing/Coding Section. References updated. Medical Director review 8/2021. (sk)
- 9/13/22 Reviewed by Avalon 2nd Quarter 2022 CAB. Scientific Background updated. Table of Terminology added. When covered section updated with two additional allowable indications. Policy Guidelines updated. References updated. Medical Director review 8/2022. (sk)

### **For Policy Titled Human Immunodeficiency Virus**

- 5/30/23 Reviewed by Avalon 1<sup>st</sup> Quarter 2023 CAB. Policies titled Plasma HIV-1 and HIV-2 RNA Quantification for HIV Infection” and “HIV Genotyping and Phenotyping” combined to form new policy titled “Human Immunodeficiency Virus”. Scientific Background updated. Table of Terminology updated. Policy Guidelines updated. References updated and added. Codes 86689, 86701, 86702, 86703, 87389, 87390, 87391, 87534, 87535, 87537, 87538, 87806, 87900, 0219U, G0432, G0433, G0435, G0475, and S3645 added to Billing/Coding section. Medical Director review 4/2023. (sk)
- 6/30/23 Added “For Policy Titled Human Immunodeficiency Virus” to Policy Implementation Section for clarity. (ldh)

---

Medical policy is not an authorization, certification, explanation of benefits or a contract. Benefits and eligibility are determined before medical guidelines and payment guidelines are applied. Benefits are determined by the group contract and subscriber certificate that is in effect at the time services are rendered. This document is solely provided for informational purposes only and is based on research of current medical literature and review of common medical practices in the treatment and diagnosis of disease. Medical practices and knowledge are constantly changing and BCBSNC reserves the right to review and revise its medical policies periodically.